

Sleep as a Fundamental Pillar of Health

What is Sleep?

Sleep is a natural state of reduced activity which marked by dynamic changes in brain wave activity, breathing patterns, heart rate, body temperature, and other physiological functions. These tightly regulated changes reflect an active and coordinated process in the brain, rather than a passive shutdown—providing the basis for understanding sleep-wake regulation.

Sleep-Wake Regulation

The sleep-wake system is thought to be regulated by the interplay of two major processes, one that promotes sleep (process S) and one that maintains wakefulness (process C).

Two primary processes govern this regulation:

1. Homeostatic Sleep Drive (Process S)
 - Increases progressively with prolonged wakefulness
 - Creates pressure to initiate sleep
 - Dissipates during sleep, particularly during deep NREM sleep
2. Circadian Rhythm (Process C)
 - Controlled by the suprachiasmatic nucleus (SCN) in the hypothalamus
 - Synchronizes sleep–wake timing with the 24-hour light–dark cycle
 - Regulates alertness, hormone secretion, and body temperature




Fig 1. Cicardian Rhythms and sleep-wake homeostasis

At the neurobiological level, sleep–wake transitions are explained by the flip-flop model of sleep regulation, in which sleep-promoting and wake-promoting neural circuits mutually inhibit each other. When wake-promoting neurons are active, they inhibit the VLPO and maintain alertness. Conversely, when the VLPO becomes active, it inhibits arousal systems and promotes sleep.




Fig. 3. Flip-flop switch

The flip-flop mechanism enables **rapid and stable transitions** between sleep and wake states, minimizing unstable intermediate states.

Sleep Architecture




Fig. 2. Normal sleep architecture

Sleep architecture refers to the **organized and predictable cyclical structure of sleep** that alternates between two major phases: **Non–Rapid Eye Movement (NREM) sleep** and **Rapid Eye Movement (REM) sleep**. NREM accounts for approximately 75% to 80% of total sleep, and REM accounts for the remaining 20% to 25% of sleep. On average, a night of sleep consists of 4 to 5 repetitions of this cycle.

The details of sleep architecture is below:

- NREM stage 1 (Light Sleep): Shallow stage of sleep where a person is still easily awoken. It lasts 1 to 7 minutes. Comprising ~ 5% of total sleep time
- NREM Stage 2 (Deeper Sleep): lasts approximately 10 to 25 minutes in the initial cycle of sleep but progresses to consume 45% of the total sleep cycle later in the night.
- NREM Stage 3 (Deepest Non-REM Sleep): lasts about 20 to 40 minutes, initially. This is critical stage for physical restoration and tissue repair. Accounts for 25% of total sleep time.
- REM stage: REM sleep is characterized by **rapid eye movements, vivid dreaming, and near-complete voluntary muscle paralysis**. This stage plays a key role in **memory consolidation, emotional regulation, and cognitive processing**. Despite muscle atonia, brain activity during REM sleep is high, with cerebral metabolism increasing by up to **20%** compared to wakefulness. REM sleep accounts for approximately **25%** of total sleep time and becomes progressively longer toward the morning hours.

Preservation of normal sleep architecture is essential for optimal physiological and neurological function. Disruptions to specific sleep stages—such as reduced deep NREM sleep or fragmented REM sleep—are associated with impaired recovery, cognitive dysfunction, and poor clinical outcomes, particularly in patients with insomnia or neurological disorders.

Importance of Sleep as Pillar of Health

Sleep is an active, highly regulated biological process that plays a fundamental role in maintaining brain function, physiological homeostasis, and overall health. It is governed by the interaction between homeostatic sleep drive and circadian rhythm, with sleep–wake transitions controlled at the neurobiological level by a flip-flop switch mechanism involving mutual inhibition between sleep- and wake-promoting neural circuits. This tightly coordinated system ensures stable sleep and wake states and preserves normal sleep architecture, which consists of cyclical transitions between NREM and REM sleep across the night.

Preservation of healthy sleep architecture is essential for physical restoration, cognitive performance, emotional regulation, and neural plasticity. Disruptions in sleep regulation—such as reduced deep NREM sleep, fragmented REM sleep, or unstable sleep–wake transitions—are associated with impaired recovery, cognitive dysfunction, and poorer clinical outcomes, particularly in patients with insomnia or neurological disorders. Recognizing sleep as a critical component of health underscores the importance of early identification and appropriate management of sleep disturbances as part of comprehensive patient care.

References:

1. Science of Sleep: What is Sleep. Available at <https://sleep.hms.harvard.edu/education-training/public-education/sleep-and-health-education-program/sleep-health-education-47>
2. Brinkman JE, et al. 2023. Physiology of Sleep. StatPearls. Treasure Island. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK482512/>
3. Patel, et al. 2024. Physiology, sleep stages. *StatPearls*. StatPearls Publishing, Treasure Island. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK526132/>
4. Institute of Medicine (US) Committee on Sleep Medicine and Research; Colten HR, Altevogt BM, editors. Sleep Disorders and Sleep Deprivation: An Unmet Public Health Problem. Washington (DC): National Academies Press(US); 2006. 2, Sleep Physiology. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK19956/>
5. Luthi, Anita. 2016. Sleep: Switching Off the Off-Switch. *Current Biology* 26, R756–R777.